

Investigation of the 5n and α 4n channels in the ^{20}Ne on ^{244}Pu reaction

R. Dressler¹, B. Eichler¹, D. Piguet¹, R. Eichler^{1,2}, H.W. Gäggeler^{1,2}, F. Haenssler², S. Soverna², M. Schädel³, E. Schimpf³, K. Eberhardt⁴, P. Thörle⁴, N. Trautmann⁴

¹ Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland

² Departement für Chemie und Biochemie, Universität Bern, Freiestrasse 3, CH-3012 Bern, Switzerland

³ Gesellschaft für Schwerionenforschung mbH, Planckstrasse 1, D-64291 Darmstadt, Germany

⁴ Institut für Kernchemie, Universität Mainz, Fritz-Strassmann-Weg 2, D-55128 Mainz, Germany

rugard.dressler@psi.ch

The isotope ^{259}Rf was firstly investigated by Flerov et al. [1] in the 5n channel of the reaction ^{22}Ne on ^{242}Pu . Two years later Zvara et al. [2] published first ever chemistry experiments with the element Rf ($Z=104$) based on the products of the same nuclear reaction. These experiments were several times repeated and controversially discussed by different authors in the next two decades (for a nearly complete compilation of these discussions see [3]). Flerov et al. showed in [1] in addition to the excitation function of ^{259}Rf the low energy part of the excitation function of ^{256}No , which is produced in the α 4n channel of the reaction ^{22}Ne on ^{242}Pu .

Later, only a few experiments were made to investigate nuclear reactions based on Pu and Ne [4, 5]. In our experiments we will continue the investigation of xn and α xn reactions in the region of Rf and No, which was started already in [5]. The intention is to study formation and decay properties of ^{259}Rf and ^{256}No in the reactions $^{244}\text{Pu}(^{20}\text{Ne}, 5n)$ and $^{244}\text{Pu}(^{20}\text{Ne}, \alpha 4n)$, respectively.

^{259}Rf has a half-life of (3.2 ± 0.6) s and decays with $(92 \pm 2)\%$ probability by emission of α -particles of 8.77 MeV and 8.87 MeV [6]. ^{256}No with a half-life of (2.91 ± 0.05) s decays also via emission of α -particles of 8.448 MeV and 8.402 MeV [6] with a $(99.47 \pm 0.06)\%$ branching ratio. For both nuclides a spontaneous fission decay mode was reported.

A Pu target (enriched 98.6% in ^{244}Pu) with a thickness of about 0.5 mg/cm^2 is in preparation at the University of Mainz via molecular plating of its nitrate on a Be-backing of $12 \text{ }\mu\text{m}$ thickness.

In a first experiment at the PSI Phillips cyclotron this target will be bombarded with $^{20}\text{Ne}^{6+}$, at a beam energy of 113 MeV.

The recoiling reaction products are swept out of the recoil chamber using a He-carbon-gas-jet and transported to the PSI Tape System [7] within 3 s. The aerosols are collected by impaction in vacuum on the magnetic tape during 2 s and subsequently the samples are moved in front of 8 consecutive α -PIPS-detectors (450 mm^2 active area). For the event-by-event recording the data acquisition system described in [8, 9] will be used.

In Figure 1 the excitation function given in [1] is compared with model calculations using the HIVAP code [10, 11]. Under the described experimental conditions about 15 events of ^{259}Rf at a cross-section of 0.6 nb will be expected within an 8 day irradiation at beam intensities of about $1.2 \text{ e}\mu\text{A}$. The use of the PSI Tape System inhibits the accumulation of long-lived spontaneously fissioning nuclides, such as ^{256}Fm , which are produced in transfer channels of the nuclear reaction. But the outcome of the experiment also depends very much on the purity of the target material, since by-products (e.g. $^{211-214}\text{Po}$) from nuclear reactions with heavy metal impurities in the target may disturb the unambiguous identification of ^{259}Rf and ^{256}No . In that case a fast chemical separation as described in [12] will be necessary.

The results of this first experiment, which is conducted in August 2003, will be presented.




Figure 1. The cross section of ^{259}Rf in the reaction $^{242}\text{Pu}(^{22}\text{Ne}, 5n)$. Symbols: experimental data from [1]; dotted line: model calculations using HIVAP for the reaction $^{244}\text{Pu}(^{20}\text{Ne}, 5n)$.

References

- [1] G.N. Flerov *et al.* Phys. Lett. **13**, 73 (1964).
- [2] I. Zvara *et al.* Sov. Atom. Energy **21**, 709 (1966).
- [3] E.K. Hyde *et al.* Radiochim. Acta **42**, 57 (1987).
- [4] Yu.A. Lazarev *et al.* Phys. Rev. **C62**, 064307 (2000).
- [5] R. Dressler, PhD Thesis, Univ. Bern, Switzerland (1999).
- [6] A. Artana-Cohen, Nuclear Data Sheets **88**, 275 (1999).
- [7] D.T. Jost, D. Vermeulen, IEEE Trans Nucl. Sci. **NS-36**, 1630 (1992).
- [8] R. Dressler, PSI Scientific Report 2002, Volume I 68 (2003).
- [9] S. Ritt, P.A. Amaudruz, "A versatile and compact data acquisition system MIDAS", (<http://midas.psi.ch>).
- [10] W. Reisdorf, M. Schädel, Z. Phys. **A343**, 47 (1992).
- [11] R. Dressler, PSI Scientific Report 1998, Volume I 131 (1999).
- [12] A. Türler *et al.* J. Alloys and Compounds **271**, 287 (1998).